

**Math 194**  
**Clicker Questions**

UNIT 1: SYSTEMS OF LINEAR EQUATIONS

1. Suppose you have a system of linear equations consisting of three equations in two unknowns. How many solutions are possible for such a system? *Mark all that apply.*

- (a) No solutions.  
(b) Exactly one solution.  
(c) Infinitely many solutions.

2. Consider the following system of linear equations.

$$\begin{array}{ccccrcrcrcr} x & + & 3y & + & z & = & 3 \\ 2x & + & 2y & + & 4z & = & -4 \\ 3x & + & y & - & z & = & 9 \end{array}$$

The corresponding augmented matrix is

$$\left[ \begin{array}{cccc|c} 1 & 3 & 1 & 3 & 3 \\ 2 & 2 & 4 & -4 & -4 \\ 3 & 1 & -1 & 9 & 9 \end{array} \right].$$

Which of the following augmented matrices correspond to a system of linear equations with the same solution set? *Mark all that apply.*

(a) Multiply the first row by 2:

$$\left[ \begin{array}{cccc|c} 2 & 6 & 2 & 6 & 6 \\ 2 & 2 & 4 & -4 & -4 \\ 3 & 1 & -1 & 9 & 9 \end{array} \right]$$

(b) Multiply the first column by 2:

$$\left[ \begin{array}{cccc|c} 2 & 3 & 1 & 3 & 3 \\ 4 & 2 & 4 & -4 & -4 \\ 6 & 1 & -1 & 9 & 9 \end{array} \right]$$

(c) Add the first row to the third row:

$$\left[ \begin{array}{cccc|c} 1 & 3 & 1 & 3 & 3 \\ 2 & 2 & 4 & -4 & -4 \\ 4 & 4 & 0 & 12 & 12 \end{array} \right]$$

(d) Interchange the first and second rows:

$$\left[ \begin{array}{cccc|c} 2 & 2 & 4 & -4 & -4 \\ 1 & 3 & 1 & 3 & 3 \\ 3 & 1 & -1 & 9 & 9 \end{array} \right]$$

(e) Interchange the first and second columns:

$$\left[ \begin{array}{cccc|c} 3 & 1 & 1 & 3 & 3 \\ 2 & 2 & 4 & -4 & -4 \\ 1 & 3 & -1 & 9 & 9 \end{array} \right]$$

3. Suppose the augmented matrix for a system of three linear equations in four unknowns has three pivot columns. True or False: The system must be consistent.
4. Suppose the augmented matrix for a system of three linear equations in three unknowns has a pivot in each of the first three columns. True or False: The system has exactly one solution.
5. Suppose you row-reduce the augmented matrix for a system of four linear equations in three unknowns and end up with a row of all zeros. How many solutions does the system have? *Mark all that are possible.*

- (a) No solutions  
 (b) Exactly one solution  
 (c) Infinitely many solutions

6. Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 4 \end{bmatrix}$  and  $\mathbf{v}_2 = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ . Geometrically, what is the set of *all* linear combinations of  $\{\mathbf{v}_1, \mathbf{v}_2\}$ ?

- (a) A point  
 (b) A line  
 (c) All of  $\mathcal{R}^2$

7. Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} 6 \\ 0 \\ -2 \end{bmatrix}$ . Which of the following is *not* a linear combination of  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ ?

(a)  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

(b)  $\begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix}$

(c)  $\begin{bmatrix} 3 \\ 3 \\ 6 \end{bmatrix}$

- (d) All of these are linear combinations of  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ .

8. Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} 6 \\ 0 \\ -2 \end{bmatrix}$ . Geometrically, what is the set of *all* linear combinations of  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ ?

- (a) A point  
 (b) A line  
 (c) A plane  
 (d) All of  $\mathbb{R}^3$

9. Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}$ ,  $\mathbf{v}_3 = \begin{bmatrix} 6 \\ 0 \\ -2 \end{bmatrix}$ . Which of the following sets has the same span as the set of all three vectors  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ ? *Mark all that apply.*

- (a)  $\{\mathbf{v}_1, \mathbf{v}_2\}$   
 (b)  $\{\mathbf{v}_2, \mathbf{v}_3\}$

- (c)  $\{\mathbf{v}_1, \mathbf{v}_3\}$   
 (d) None of the above
10. Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 3 \\ 0 \\ -1 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} -1 \\ 2 \\ 5 \end{bmatrix}$ . Geometrically, what is the subspace spanned by  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ ?
- (a) A point  
 (b) A line  
 (c) A plane  
 (d) All of  $\mathbb{R}^3$
11. **Theorem:** Let  $A$  be an  $m \times n$  matrix. Then the following statements are logically equivalent. That is, for a particular  $A$ , either they are all true statements or they are all false statements.
- (a) For each  $\mathbf{b}$  in  $\mathbb{R}^m$ , the equation  $A\mathbf{x} = \mathbf{b}$  has a solution.  
 (b) Each  $\mathbf{b}$  in  $\mathbb{R}^m$  is a linear combination of the columns of  $A$ .  
 (c) The columns of  $A$  span  $\mathbb{R}^m$ .  
 (d)  $A$  has a pivot position on every row.

If the four statements in this theorem are all true, what does that imply about the relative size of  $m$  and  $n$ ?

- (a)  $m \geq n$   
 (b)  $m \leq n$   
 (c)  $m = n$   
 (d) Nothing.  $m$  and  $n$  can be any positive integers.

## UNIT 2: MATRIX ALGEBRA

1. Let  $A = \begin{bmatrix} 4 & 7 & 2 \\ -1 & 0 & 8 \\ 3 & 5 & 1 \end{bmatrix}$ . The entry  $a_{2,3}$  is which of the following?
- (a) 0  
 (b) 1  
 (c) 8  
 (d) 5
2. Let  $A = \begin{bmatrix} 4 & 7 & 2 \\ -1 & 0 & 8 \\ 3 & 5 & 1 \end{bmatrix}$ . Which of the following is equal to  $\mathbf{a}_2$ ?
- (a)  $[-1 \ 0 \ 8]$   
 (b)  $\begin{bmatrix} 7 \\ 0 \\ 5 \end{bmatrix}$   
 (c) 0  
 (d) -1
3. Let  $A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$ . Which of the following is equal to  $5A$ ?

(a)  $\begin{bmatrix} 10 & 3 \\ 1 & 4 \end{bmatrix}$

(b)  $\begin{bmatrix} 10 & 15 \\ 1 & 4 \end{bmatrix}$

(c)  $\begin{bmatrix} 10 & 15 \\ 5 & 20 \end{bmatrix}$

(d)  $\begin{bmatrix} 10 & 3 \\ 1 & 20 \end{bmatrix}$

4. Let  $B = \begin{bmatrix} 4 & 7 \\ -1 & 0 \\ 5 & 3 \end{bmatrix}$ . Which of the following is equal to  $B^T$ ?

(a)  $\begin{bmatrix} 4 & -1 & 5 \\ 7 & 0 & 3 \end{bmatrix}$

(b)  $\begin{bmatrix} 7 & 4 \\ 0 & -1 \\ 3 & 5 \end{bmatrix}$

(c)  $\begin{bmatrix} 5 & 3 \\ -1 & 0 \\ 4 & 7 \end{bmatrix}$

(d)  $B^T$  cannot be computed.

5. Let  $A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$  and  $B = \begin{bmatrix} -1 & 7 \\ 0 & 2 \end{bmatrix}$ . Which of the following is equal to  $AB$ ?

(a)  $\begin{bmatrix} -2 & 21 \\ 0 & 8 \end{bmatrix}$

(b)  $\begin{bmatrix} -2 & 20 \\ -1 & 15 \end{bmatrix}$

(c)  $\begin{bmatrix} 5 & 25 \\ 2 & 8 \end{bmatrix}$

(d)  $AB$  cannot be computed.

6. If  $A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 & 0 & 7 \\ 2 & 6 & 4 \end{bmatrix}$ , and  $C = \begin{bmatrix} 9 & 8 & 7 \\ 6 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$ , then which of the following can be computed?

(Mark all that apply.)

(a)  $AB$

(b)  $BA$

(c)  $BC$

(d)  $CB$

(e)  $AC$

7. If  $A$  is an  $m \times n$  matrix and  $B$  is an  $n \times p$  matrix, then  $A\mathbf{b}_1$  is described by which of the following?

(a) The first row of  $AB$

(b) The first column of  $AB$

(c) The 1, 1-entry of  $AB$

(d) The first row of  $BA$

8. Suppose you have a simple board game called Quad that consists of four spaces, A, B, C, and D. Movement is as follows:
- (a) If you're on space  $A$ , you have a 50% chance of moving to space  $B$  and a 50% chance of staying on space  $A$ .
  - (b) If you're on space  $B$ , you have a 50% chance of moving to space  $A$  and a 50% chance of staying on space  $B$ .
  - (c) If you're on space  $C$ , you have a 30% chance of moving to space  $D$  and a 70% chance of staying on space  $C$ .
  - (d) If you're on space  $D$ , you have a 30% chance of moving to space  $C$  and a 70% chance of staying on space  $D$ .


Is the transition matrix  $P$  for this system regular? That is, is there some positive integer power  $k$  such that  $P^k$  has all positive entries?

- (a) Yes
  - (b) No
9. True or False: If  $A$  and  $B$  are square matrices with the same dimensions, then  $(A + B)(B + A) = A^2 + 2AB + B^2$ .
- (a) True
  - (b) False
10. True or False: If  $AB = AC$ , then  $B = C$ .
- (a) True
  - (b) False
11. True or False: If  $AD = 0$ , then either  $A = 0$  or  $D = 0$ .
- (a) True
  - (b) False
12. If  $A$  and  $B$  are invertible matrices, then what is  $(AB)^{-1}$ ?
- (a)  $A^{-1}B^{-1}$
  - (b)  $B^{-1}A^{-1}$
  - (c)  $AB^{-1}$
  - (d)  $BA^{-1}$
  - (e) More than one of the above is true.
  - (f) Just because  $A$  and  $B$  have inverses, this doesn't mean that  $AB$  has an inverse.
13. Which of the following matrices does not have an inverse?
- (a)  $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$
  - (b)  $\begin{bmatrix} 2 & 2 \\ 4 & 4 \end{bmatrix}$
  - (c)  $\begin{bmatrix} -1 & 0 \\ 0 & 3 \end{bmatrix}$
  - (d)  $\begin{bmatrix} 0 & 4 \\ 2 & 0 \end{bmatrix}$

- (e) All have inverses.
14. Suppose that for a square coefficient matrix  $A$ , the matrix equation  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution  $\mathbf{x} = \mathbf{0}$ . This means that
- Matrix  $A$  has no inverse.
  - Matrix  $A$  has an inverse.
  - This tells us nothing about whether  $A$  has an inverse.
15. Which of the following is a set of linearly independent vectors?

- $\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 4 \\ 7 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$
- $\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 4 \\ 7 \end{bmatrix}, \begin{bmatrix} 5 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 10 \end{bmatrix}$
- $\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 4 \\ 7 \end{bmatrix}, \begin{bmatrix} 2 \\ -2 \\ 6 \end{bmatrix}$
- $\begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} -2 \\ 4 \\ 7 \end{bmatrix}, \begin{bmatrix} 6 \\ -12 \\ -21 \end{bmatrix}$

16. Which subsets of the set of vectors shown are linearly dependent? (Mark all that apply.)



- $\mathbf{u}, \mathbf{w}$
  - $\mathbf{t}, \mathbf{w}$
  - $\mathbf{t}, \mathbf{v}$
  - $\mathbf{t}, \mathbf{u}, \mathbf{v}$
17. Suppose you wish to determine whether a set of vectors  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$  is linearly independent. You form the matrix  $A = [\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3 \ \mathbf{v}_4]$ , and you calculate its reduced row echelon form,  $R = \begin{bmatrix} 1 & 0 & 2 & 1 \\ 0 & 1 & 3 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ . You now decide to write  $\mathbf{v}_2$  as a linear combination of  $\mathbf{v}_1, \mathbf{v}_3$ , and  $\mathbf{v}_4$ . Which is a correct linear combination?

- (a)  $\mathbf{v}_2 = 3\mathbf{v}_3 + \mathbf{v}_4$
- (b)  $\mathbf{v}_2 = -3\mathbf{v}_3 - \mathbf{v}_4$
- (c)  $\mathbf{v}_2 = \mathbf{v}_4 - 3\mathbf{v}_3$
- (d)  $\mathbf{v}_2 = -\mathbf{v}_1 + \mathbf{v}_4$
- (e) I don't know.

UNIT 3: SUBSPACES

1. Consider the set  $\{\mathbf{x} \in \mathbb{R}^3 \mid x_1 = 2x_3\}$ . If  $\mathbf{x}$  and  $\mathbf{y}$  are two vectors in this set, is any linear combination of  $\mathbf{x}$  and  $\mathbf{y}$  also in the set?
  - (a) Yes
  - (b) No
2. Consider the set  $\{\mathbf{x} \in \mathbb{R}^3 \mid x_1 = 2\}$ . If  $\mathbf{x}$  and  $\mathbf{y}$  are two vectors in this set, is any linear combination of  $\mathbf{x}$  and  $\mathbf{y}$  also in the set?
  - (a) Yes
  - (b) No
3. Consider the set  $\{\mathbf{x} \in \mathbb{R}^3 \mid x_1 = x_2 = x_3 = 0\}$ . Is this set a subspace of  $\mathbb{R}^3$ ?
  - (a) Yes
  - (b) No
4. Suppose  $A$  is a fixed  $3 \times 3$  matrix and  $\mathbf{b}$  is a fixed  $3 \times 1$  vector. Consider the set  $S = \{\mathbf{x} \in \mathbb{R}^3 \mid A\mathbf{x} = \mathbf{b}\}$ . Is  $S$  a subspace of  $\mathbb{R}^3$ ?
  - (a) Yes
  - (b) No
5. Suppose  $A$  is a fixed  $3 \times 3$  matrix. Consider the set  $S = \{\mathbf{x} \in \mathbb{R}^3 \mid A\mathbf{x} = \mathbf{0}\}$ . Is  $S$  a subspace of  $\mathbb{R}^3$ ?
  - (a) Yes
  - (b) No
6. Which of the following sets is a basis for  $\mathbb{R}^3$ ?
  - (a)  $\begin{bmatrix} 4 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix}, \begin{bmatrix} 7 \\ 5 \\ 0 \end{bmatrix}$
  - (b)  $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}, \begin{bmatrix} 10 \\ 0 \\ 7 \end{bmatrix}$
  - (c)  $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ -2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 3 \end{bmatrix}$
  - (d)  $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 7 \\ 1 \end{bmatrix}$
7. True or False: If  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  and  $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m\}$  are both bases for a subspace  $S$ , then  $n = m$ .
  - (a) True

(b) False

8. Let  $A = \begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 2 & -1 & 1 \\ 0 & 1 & 0 & 1 \\ 3 & -1 & 0 & 2 \end{bmatrix}$ . The reduced row echelon form of  $A$  is  $R = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ . What is the rank of  $A$ ?

- (a) 1
- (b) 2
- (c) 3
- (d) 4

9. True or False: Two row equivalent matrices have the same column space.

- (a) True
- (b) False

10. Let  $A = \begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 2 & -1 & 1 \\ 0 & 1 & 0 & 1 \\ 3 & -1 & 0 & 2 \end{bmatrix}$ . The reduced row echelon form of  $A$  is  $R = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ . What is the dimension of the column space of  $A$ ?

- (a) 1
- (b) 2
- (c) 3
- (d) 4

11. Let  $A = \begin{bmatrix} 1 & 0 & 2 & 3 \\ 0 & 2 & -1 & 1 \\ 0 & 1 & 0 & 1 \\ 3 & -1 & 0 & 2 \end{bmatrix}$ . The reduced row echelon form of  $A$  is  $R = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ . Which columns of  $A$  would form a basis for the column space of  $A$ ?

- (a) All four
- (b) The first three
- (c) Any three
- (d) Any two

12. True or False: The dimension of the row space of a matrix is equal to the dimension of the column space of that matrix.

- (a) True
- (b) False

13. True or False: If  $A$  is an  $m \times n$  matrix, then the dimension of the row space of  $A$  plus the dimension of the nullspace of  $A$  equals  $n$ .

- (a) True
- (b) False



1. Which of the following is not a linear transformation?

(a)  $L(\mathbf{x}) = \begin{bmatrix} x_1 \\ x_2 + 1 \end{bmatrix}$

(b)  $L(\mathbf{x}) = \begin{bmatrix} x_1 - 2x_2 \\ x_1 \end{bmatrix}$

(c)  $L(\mathbf{x}) = \begin{bmatrix} 4x_2 \\ x_1 - 2x_2 \end{bmatrix}$

(d)  $L(\mathbf{x}) = \begin{bmatrix} x_1 \\ 0 \end{bmatrix}$

(e) All are linear transformations

(f) None are linear transformations

2. Consider the linear transformation  $L : \mathbb{R}^4 \Rightarrow \mathbb{R}^3$  defined by  $L(\mathbf{x}) = A\mathbf{x}$  where  $A = \begin{bmatrix} 1 & -4 & 8 & 1 \\ 0 & 2 & -1 & 3 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ .

Which of the following is true?

(a)  $L$  is onto, but not one-to-one.

(b)  $L$  is one-to-one, but not onto.

(c)  $L$  is both onto and one-to-one.

(d)  $L$  is neither onto nor one-to-one.

3. Consider the linear transformation  $L : \mathbb{R}^2 \Rightarrow \mathbb{R}^3$  defined by  $L(\mathbf{x}) = A\mathbf{x}$  where  $A = \begin{bmatrix} 3 & 1 \\ 5 & 7 \\ 1 & 3 \end{bmatrix}$ . Which of the following is true?

(a)  $L$  is onto, but not one-to-one.

(b)  $L$  is one-to-one, but not onto.

(c)  $L$  is both onto and one-to-one.

(d)  $L$  is neither onto nor one-to-one.

4. Suppose you want to translate the vector  $\begin{bmatrix} x \\ y \end{bmatrix}$  by 3 units to the right and 4 units up in  $\mathbb{R}^2$ . Left-

multiplying  $\begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$  by which of the following matrices will yield the homogeneous coordinates of the correct image?

(a)  $\begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 4 & 0 & 1 \end{bmatrix}$

(b)  $\begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix}$


(c)  $\begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{bmatrix}$

(d)  $\begin{bmatrix} 1 & 0 & 4 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{bmatrix}$

5. Suppose  $A = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}$ . Let  $T$  be the transformation with standard matrix  $AB$ . What 2D transformation is performed by  $T$  when it acts on the homogeneous coordinates of a point  $(x, y)$ ?
- A rotation by 90 degrees clockwise, then a translation by 2 units right and 1 unit down
  - A translation by 2 units right and 1 unit down, then a rotation by 90 degrees clockwise
  - Both of the above are correct.
  - None of the above are correct.
6. Which of the following types of linear transformation is NOT invertible?
- rotations
  - reflections
  - shears
  - projections
  - translations

#### UNIT 5: EIGENVALUES AND EIGENVECTORS

1. The vector  $\mathbf{t}$  is an eigenvector of the matrix  $A$ . What could be the result of the product  $A\mathbf{t}$ ?



- $A\mathbf{t} = \mathbf{u}$
- $A\mathbf{t} = \mathbf{v}$
- $A\mathbf{t} = \mathbf{w}$
- None of the above

ANSWERS:

UNIT 1: SYSTEMS OF LINEAR EQUATIONS

1. (a), (b), and (c)
2. (a), (c), and (d)
3. False
4. True
5. (a), (b), and (c)
6. (c)
7. (a)
8. (c)
9. (a) and (c)
10. (c)
11. (b)

UNIT 2: MATRIX ALGEBRA

1. (c)
2. (b)
3. (c)
4. (a)
5. (b)
6. (a) and (c)
7. (b)
8. No
9. False
10. False
11. False
12. (b)
13. (b)
14. (b)
15. (c)
16. (b) and (d)
17. (d)

### UNIT 3: SUBSPACES

1. (a)
2. (b)
3. (a)
4. (b)
5. (a)
6. (c)
7. True
8. (c)
9. False
10. (c)
11. (c)
12. True
13. True

### UNIT 4: LINEAR TRANSFORMATIONS

1. (a)
2. (a)
3. (b)
4. (c)
5. (b)
6. (d)

### UNIT 5: EIGENVALUES AND EIGENVECTORS

1. (c)